

112 x 180mm

Sofac-P 50mg
Tablets

(Diclofenac Potassium USP)

سوفیک-پی
۵۰ ملی گرام
ٹیبلیٹس

(ڈیکلو فینیک پوٹاشیم)

Description:

Sofac-P, the brand of Diclofenac Potassium, is an antiinflammatory, analgesic and antipyretic agent.

Composition:

Each film coated tablet contains:
Diclofenac Potassium USP.....50mg.

Action:

1. **Sofac-P** is a nonsteroidal anti-inflammatory agent that possesses analgesic and antipyretic activities. Its mode of action, like that of other nonsteroidal anti-inflammatory agents, is not known; however, its therapeutic action is not due to pituitary-adrenal stimulation, but may be from the inhibitory action in the synthesis of prostaglandin by inhibiting enzyme Cyclo-oxygenase.
2. **Sofac-P** exhibits very marked antirheumatic, anti-inflammatory, analgesic and antipyretic properties and is thus suitable for use in inflammatory and degenerative rheumatic diseases, as well as for the treatment of post-traumatic inflammation and swelling.

Indications:

Inflammatory and degenerative forms of rheumatism, juvenile rheumatoid arthritis, ankylosing spondylitis, osteoarthritis and spondyl spondyl arthritis.
- Painful syndromes of the vertebral column.
- Non-Article rheumatism.
- Acute attack of gout.
- painful post traumatic and post operative inflammation and swelling of orthopaedic surgery.
- painful and inflammatory conditions in gynaecology, e.g. Primary dysmenorrhoea and adnexitis.
- As an adjuvant therapy in severe painful inflammatory infection of ear, nose and throat.
- painful post-operative, inflammatory condition of dental surgery.

Dosage & Administration:

Adults:

Initial recommended dose is 100 to 150mg per day but in mild cases and long duration of treatment the usual dose may be 75 to 100mg per day. Dose should be divided into two or three parts in a day. Tablet should be taken with meal or after meal with water.

Children:

The use of **Sofac-P** is not recommended in children below 14 years of age.

Side Effects:

At the start of treatment, patient may complain of epigastric pain, eructation, nausea and diarrhoea, headache, or dizziness. These side effects, however, are usually of a mild nature and generally disappear after a few days despite continuation of the medication. Skin rashes may also occur very occasionally.

Precautions:

1. In accordance with currently accepted views on the employment of drugs during pregnancy, **Sofac-P** should not be prescribed in the first three months of pregnancy unless its use is strictly indicated.
2. Patients with gastro-intestinal disturbances or with history of peptic ulcer, as well as those suffering from severe hepatic or renal damage, should be kept under close observation.

Dosage: As directed by the physician.

Instructions:

Store below 30°C in a cool dry place & protect from light.

Keep out of the reach of children.

To be sold on the prescription of a registered medical practitioner only.

Presentation:

Sofac-P tablet supplied in 2 x 10's in blister pack.

Saffron

Pharmaceuticals (Pvt.) Ltd.

19 Km. Sheikhpura Road, Faisalabad-Pakistan.

Tel: +92-41-2423914-18, Fax: +92-41-2423919

E-mail: info@saffronpharma.com

www.saffronpharma.com

خوراک:

ڈاکٹر کی ہدایت کے مطابق استعمال کریں۔

ہدایات:

دوا کو ۳۰°C سے کم درجہ حرارت پر

روشنی سے بچا کر خشک جگہ پر رکھیں۔ بچوں کی پہنچ سے دور رکھیں۔

صرف رجسٹرڈ ڈاکٹر کے نسخے پر ہی فروخت کریں۔

QAR No. 1012005